



Ice Sheet System model

Ice flow models


Eric LAROUR¹, Eric RIGNOT^{1,3}, Mathieu MORLIGHEM^{1,2}, Hélène SEROUSSI^{1,2} Chris BORSTAD¹, Feras HABBAL^{1,3}, Daria HALKIDES^{1,4}, Behnaz KHAKBAZ¹, John SCHIERMEIER¹, Nicole SCHLEGEL¹

¹Jet Propulsion Laboratory - California Institute of Technology

²Laboratoire MSSMat, École Centrale Paris, France

³University of California, Irvine

⁴Joint Institute for Regional Earth System Science & Engineering, UCLA



Ice flow models

Larour et al.

Ice flow equations

- Approximations implemented
- Ice flow equation
- Diagnostic parameters
- Boundary conditions

Combining models

- Methods implemented in ISSM
- Penalties
- Tiling method
- Utilization

Outline

① Ice flow equations

- Approximations implemented
- Ice flow equation
- Diagnostic parameters
- Boundary conditions

② Combining models

- Methods implemented in ISSM
- Penalties
- Tiling method
- Utilization

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters
Boundary conditions

Combining models

Methods Implemented in
ISSM

Penalties

Tiling method

Utilization

Ice Sheet flow equations

Incompressibility

$$\forall \mathbf{x} \in \Omega \quad \nabla \cdot \mathbf{v} = \text{Tr}(\dot{\boldsymbol{\epsilon}}) = \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0 \quad (1)$$

- $\mathbf{v} = (u, v, w)$ ice velocity (m/yr)
- $\dot{\boldsymbol{\epsilon}}$ strain rate tensor (yr^{-1})

Incompressible viscous fluid

$$\sigma' = 2\mu\dot{\boldsymbol{\epsilon}} \quad (2)$$

- σ' deviatoric stress
- μ ice viscosity
- $\dot{\boldsymbol{\epsilon}}$ strain rate tensor

Glen's flow law

$$\mu = \frac{B}{2\dot{\epsilon}_e^{\frac{n-1}{n}}} \quad (3)$$

- B ice hardness
- n Glen's law coefficient ($n = 3$)
- $\dot{\epsilon}_e$ effective strain rate (second invariant)



Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods Implemented in ISSM

Penalties

Tiling method

Utilization

Ice Sheet flow equations

Conservation of momentum

$$\forall \mathbf{x} \in \Omega \quad \nabla \cdot \boldsymbol{\sigma}' - \nabla P + \rho \mathbf{g} = \mathbf{0} \quad (4)$$

Assumptions:

- ① Stokes flow (quasi-static assumption)
- ② Coriolis effect negligible

Boundary conditions

Ice/Air interface: Free surface

$$\Gamma_s \quad \boldsymbol{\sigma} \cdot \mathbf{n} = P_{atm} \quad \mathbf{n} \simeq \mathbf{0}$$

Ice/Ocean interface: water pressure

$$\Gamma_w \quad \boldsymbol{\sigma} \cdot \mathbf{n} = P_w \quad \mathbf{n}$$

Ice/Bedrock interface (1): lateral friction

$$\Gamma_b \quad (\boldsymbol{\sigma} \cdot \mathbf{n} + \beta \mathbf{v})_{\parallel} = \mathbf{0}$$

Ice/Bedrock interface (2): impenetrability

$$\Gamma_b \quad \mathbf{v} \cdot \mathbf{n} = \mathbf{0}$$

Side boundaries: Dirichlet

$$\Gamma_u \quad \mathbf{v} = \mathbf{v}_{obs}$$

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters
Boundary conditions

Combining models

Methods implemented in
ISSM

Penalties


Tiling method

Utilization

Models description

Full-Stokes model:

- Momentum balance + incompressibility
- 3D model
- Four unknowns (v_x, v_y, v_z, p)



Model equations

$$\begin{cases} \frac{\partial}{\partial x} \left(2\mu \frac{\partial v_x}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) - \frac{\partial p}{\partial x} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial y} \left(2\mu \frac{\partial v_y}{\partial y} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) - \frac{\partial p}{\partial y} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) + \frac{\partial}{\partial z} \left(2\mu \frac{\partial v_z}{\partial z} \right) - \frac{\partial p}{\partial z} - \rho g = 0 \\ \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} = 0 \end{cases}$$

JPL**ISSM**

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods Implemented in ISSM

Penalties

Tiling method


Utilization

Models description

Blatter-Pattyn (BP)

Higher-order model:

- [Blatter, 1995, Pattyn, 2003]
- 3D model
- Horizontal and vertical velocity decoupled
- $2(v_x, v_y) + 1(v_z)$ unknowns

 (v_x, v_y)

Model equations

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x} \left(2\mu \frac{\partial v_x}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) - \frac{\partial p}{\partial x} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial y} \left(2\mu \frac{\partial v_y}{\partial y} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) - \frac{\partial p}{\partial y} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) + \frac{\partial}{\partial z} \left(2\mu \frac{\partial v_z}{\partial z} \right) - \frac{\partial p}{\partial z} - \rho g = 0 \end{array} \right.$$

$$\frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} = 0$$

JPL

ISSM

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods Implemented in ISSM

Penalties

Tiling method


Utilization

Models description

MacAyeal-Morland (SSA)

Shelfy-stream approximation:

- [MacAyeal, 1989]
- 2D model
- Horizontal and vertical velocity decoupled
- $2(v_x, v_y) + 1(v_z)$ unknowns



Model equations

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x} \left(2\mu \frac{\partial v_x}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) - \frac{\partial p}{\partial x} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial y} \left(2\mu \frac{\partial v_y}{\partial y} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) - \frac{\partial p}{\partial y} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) + \frac{\partial}{\partial z} \left(2\mu \frac{\partial v_z}{\partial z} \right) - \frac{\partial p}{\partial z} - \rho g = 0 \\ \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} = 0 \end{array} \right.$$

JPL**ISSM**

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods implemented in ISSM

Penalties

Tiling method


Utilization

Models description

Hutter (SIA)

Shallow ice approximation:

- [Hutter, 1983]
- 3D analytical model
- 2 unknowns (v_x, v_y) computed separately

 (v_x, v_y)

Model equations

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x} \left(2\mu \frac{\partial v_x}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) - \frac{\partial p}{\partial x} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial y} \left(2\mu \frac{\partial v_y}{\partial y} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) - \frac{\partial p}{\partial y} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) + \frac{\partial}{\partial z} \left(2\mu \frac{\partial v_z}{\partial z} \right) - \frac{\partial p}{\partial z} - \rho g = 0 \\ \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} = 0 \end{array} \right.$$

JPL

ISSM

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods implemented in ISSM

Penalties

Tiling method

Utilization

Material non-linearity

Model equations

$$\begin{cases} \frac{\partial}{\partial x} \left(2\mu \frac{\partial v_x}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) - \frac{\partial p}{\partial x} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial y} + \mu \frac{\partial v_y}{\partial x} \right) + \frac{\partial}{\partial y} \left(2\mu \frac{\partial v_y}{\partial y} \right) + \frac{\partial}{\partial z} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) - \frac{\partial p}{\partial y} = 0 \\ \frac{\partial}{\partial x} \left(\mu \frac{\partial v_x}{\partial z} + \mu \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial y} \left(\mu \frac{\partial v_y}{\partial z} + \mu \frac{\partial v_z}{\partial y} \right) + \frac{\partial}{\partial z} \left(2\mu \frac{\partial v_z}{\partial z} \right) - \frac{\partial p}{\partial z} - \rho g = 0 \\ \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} = 0 \end{cases}$$

Glen's flow law

$$\mu = \frac{B}{2 \varepsilon_e^{\frac{n-1}{n}}} \quad (5)$$

- B ice hardness
- n Glen's law coefficient ($n = 3$)
- $\dot{\varepsilon}_e$ effective strain rate (second invariant)

→ Treatment of non-linearity with fixed point



Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods Implemented in ISSM


Penalties

Tiling method

Utilization

Material non-linearity

Treatment of non-linearity with fixed point:



Vertical velocity computed with incompressibility for 2d shelfy-stream and 3d Blatter/Pattyn modes.

[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)

Approximations implemented

[Ice flow equation](#)Diagnostic parameters
Boundary conditions[Combining models](#)Methods Implemented in
ISSM

Penalties

Tiling method

Utilization

Flow equation

`setflowequation` is used to generate the approximation used to compute the velocity

- Arguments:
 - ① model
 - ② approximation names
 - ③ approximation domains
- Domains can be Argus files or array of element flags
- Approximation available
 - stokes (Full-Stokes model)
 - pattyn (Higher-order model)
 - macayeal (Shallow Shelf Approximation)
 - hutter (Shallow Ice Approximation)
- Possibility of coupling models

[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)[Approximations implemented](#)[Ice flow equation](#)[Diagnostic parameters
Boundary conditions](#)[Combining models](#)[Methods Implemented in
ISSM](#)[Penalties](#)[Tiling method](#)[Utilization](#)

Flow equation

`setflowequation` is used to generate the approximation used to compute the velocity

- Examples

```
1 md=setflowequation(md,'hutter','all')
2 md=setflowequation(md,'stokes','all')
3 md=setflowequation(md,'macayeal','all')
4 md=setflowequation(md,'pattyn','all')
```

- To display the type of approximation:


```
1 >> plotmodel(md,'data','elements_type')
```

[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)[Approximations implemented](#)[Ice flow equation](#)[Diagnostic parameters](#)[Boundary conditions](#)[Combining models](#)[Methods implemented in ISSM](#)[Penalties](#)[Tiling method](#)[Utilization](#)

Flow equation

- To display the type of approximation:

```
1 >> plotmodel(md,'data','elements_type')
```



[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)[Approximations implemented](#)[Ice flow equation](#)[Diagnostic parameters](#)
[Boundary conditions](#)[Combining models](#)[Methods Implemented in ISSM](#)[Penalties](#)[Tiling method](#)[Utilization](#)

Flow equation class

```
1  >> md.flowequation
2
3  ans =
4
5  flow equation parameters:
6      ismacayealpattyn   : 0    -- is the macayeal or pattyn approximation used ?
7      isshutter           : 0    -- is the shallow ice approximation used ?
8      isstokes             : 0    -- are the Full-Stokes equations used ?
9      vertex_equation     : N/A   -- flow equation for each vertex
10     element_equation    : N/A   -- flow equation for each element
11     bordermacayeal     : N/A   -- vertices on MacAyeal's border (for tiling)
12     borderpattyn        : N/A   -- vertices on Pattyn's border (for tiling)
13     borderstokes         : N/A   -- vertices on Stokes' border (for tiling)
```

Ice flow models

Larour et al.

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary

Methods implemented in ISSM

Penalty

Tiling method

Utilization

[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)[Approximations implemented](#)[Ice flow equation](#)[Diagnostic parameters](#)[Boundary conditions](#)[Combining models](#)[Methods Implemented in ISSM](#)[Penalties](#)[Tiling method](#)[Utilization](#)

Boundary conditions

Boundary conditions created automatically or manually

- Automatically:

```
1  >> md=SetIceSheetBC(md)
2  >> md=SetIceShelfBC(md, 'Front.exp')
3  >> md=SetMarineIceSheefBC(md, 'Front.exp')
```

- Manually: fields to change

- `md.diagnostic.spcvx`
- `md.diagnostic.spcvy`
- `md.diagnostic.spcvz`
- `md.diagnostic.icefront`

- To display the boundary conditions

```
1  >> plotmodel(md, 'data', 'BC')
```

Ice flow models

Larour et al.

Ice flow equations

Approximations
implemented

Ice flow equation

Diagnostic parameters
Boundary conditions

Combining models

Methods implemented in
ISSM

Penalties


Tiling method

Utilization

Models description

"Everything should be made as simple as possible, but no simpler." Albert Einstein

Model	Dim.	Unknowns	Reference
Full-Stokes (FS)	3d	4	[Stokes, 1845]
Blatter-Pattyn (BP)	3d	2 + 1	[Blatter, 1995, Pattyn, 2003]
Shallow shelf (SSA)	2d	2 + 1	[MacAyeal, 1989]
Shallow ice (SIA)	2d	2 + 1	[Hutter, 1983]



Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters
Boundary conditions

Combining models

Methods Implemented in
ISSM



Penalties

Tiling method

Utilization

Penalty method

- Only to couple SSA and HO
- Very stiff spring to penalize differences between degrees of freedom



Using penalties to couple models:

```
1 md=setflowequation(md,'macayeal','FloatingIce.exp','fill','pattyn','coupling','penalties')
```

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters
Boundary conditions

Combining models

Methods Implemented in
ISSM


Penalties

Tiling method

Utilization

Domain Decomposition

- $\Omega = \Omega_1 \cup \Omega_2$
- $\Omega_S = \Omega_1 \cap \Omega_2 \neq \emptyset$
- $\mathbf{u} = \mathbf{u}_1|_{\Omega_1} + \mathbf{u}_2|_{\Omega_2} \in \tilde{V}(\Omega) = (\mathcal{V}_1(\Omega_1) + \mathcal{V}_2(\Omega_2))$



Find $\mathbf{u} = \mathbf{u}_1|_{\Omega_1} + \mathbf{u}_2|_{\Omega_2} \in \tilde{V}$,

$$\forall (\mathbf{v}_1, \mathbf{v}_2) \in \tilde{V} \quad a(\mathbf{u}_1 + \mathbf{u}_2, \mathbf{v}_1 + \mathbf{v}_2) = l(\mathbf{v}_1 + \mathbf{v}_2)$$

→ Infinite number of solutions for the continuous problem

JPL

ISSM

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models


Methods Implemented in ISSM

Penalties

Tiling method


Utilization

Discretization



We take advantage of the discretization to avoid the redundancy:

- Create one layer of elements in the superposition zone



Ice flow models

Larour et al.

Ice flow equations

Approximations

implemented

ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods implemented in
ISSM

Penalties

Tiling method

Utilization

Multi-model formulation

Two different models: a_1 , a_2 and l_1 , l_2 Find $\mathbf{u} = \mathbf{u}_1|_{\Omega_1} + \mathbf{u}_2|_{\Omega_2} \in (\mathcal{V}_1 + \mathcal{V}_2)$, such that:

$$\forall \mathbf{v} = \mathbf{v}_1|_{\Omega_1} + \mathbf{v}_2|_{\Omega_2} \in (\mathcal{V}_1 + \mathcal{V}_2)$$

$$\underbrace{a_1(\mathbf{u}_1|_{\Omega_1}, \mathbf{v}_1|_{\Omega_1})}_{\text{model 1}} + \underbrace{a_2(\mathbf{u}_2|_{\Omega_2}, \mathbf{v}_2|_{\Omega_2})}_{\text{model 2}} + \\ \underbrace{a_2(\mathbf{u}_1|_{\Omega_1}, \mathbf{v}_2|_{\Omega_2}) + a_1(\mathbf{u}_2|_{\Omega_2}, \mathbf{v}_1|_{\Omega_1})}_{\text{model coupling}} \\ = l_1(\mathbf{v}_1|_{\Omega_1}) + l_2(\mathbf{v}_2|_{\Omega_2})$$

- Coupling different mechanical models
- Easy to implement (local modification of stiffness matrices)

[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)[Approximations implemented](#)[Ice flow equation](#)[Diagnostic parameters](#)[Boundary conditions](#)[Combining models](#)[Methods implemented in ISSM](#)[Penalties](#)[Tiling method](#)[Utilization](#)

Flow equation

`setflowequation` is used to generate the approximation used to compute the velocity

- Examples

```
1 md=setflowequation(md,'pattyn',md.elementongroundedice,'fill','macayeal','coupling','penalties')
2 md=setflowequation(md,'pattyn',md.elementongroundedice,'fill','macayeal','coupling','tiling')
3 md=setflowequation(md,'stokes','Contour.exp','fill','pattyn')
```

- Use `exptool` to create EXP contours

```
1 >> exptool('Contour.exp')
```

Ice flow models

Larour et al.

Ice flow equations

Approximations implemented

Ice flow equation

Diagnostic parameters

Boundary conditions

Combining models

Methods Implemented in ISSM

Penalties


Tiling method

Utilization

Flow equation

- To display the type of approximation:

```
1  >> ...
plotmodel(md,'data','elements_type','edgecolor','k','expdisp','Contour.exp')
```



[Ice flow models](#)[Larour et al.](#)[Ice flow equations](#)[Approximations implemented](#)[Ice flow equation](#)[Diagnostic parameters
Boundary conditions](#)[Combining models](#)[Methods Implemented in ISSM](#)[Penalties](#)[Tiling method](#)[Utilization](#)

Bibliography I

**Blatter, H. (1995).**

Velocity and stress-fields in grounded glaciers: A simple algorithm for including deviatoric stress gradients.

J. Glaciol., 41(138):333–344.

**Hutter, K. (1983).**

Theoretical glaciology: material science of ice and the mechanics of glaciers and ice sheets.

D. Reidel Publishing Co, Dordrecht, The Netherlands.

**MacAyeal, D. (1989).**

Large-scale ice flow over a viscous basal sediment - Theory and application to ice stream-B, Antarctica.

J. Geophys. Res., 94(B4):4071–4087.

**Pattyn, F. (2003).**

A new three-dimensional higher-order thermomechanical ice sheet model: Basic sensitivity, ice stream development, and ice flow across subglacial lakes.

J. Geophys. Res., 108(B8):1–15.

**Stokes, G. (1845).**

On the theories of internal friction of fluids in motion.

Trans. Cambridge Philos. Soc., 8:287–305.

A wide-angle photograph of a desolate, cold landscape, likely an Antarctic or Arctic scene. The foreground is a flat, light blue-grey expanse of sea ice or snow. In the middle ground, a range of majestic, snow-capped mountains rises against a clear, pale blue sky. The mountains have rugged peaks and deep, shadowed valleys. A single, bold, black text "Thanks!" is centered in the upper portion of the image.

Thanks!